

Relativistic Effects on Neutrino Pair Annihilation above a Kerr Black Hole with the Accretion Disk

Katsuaki Asano¹ and Takeshi Fukuyama²

¹Department of Earth and Space Science, Graduate School of Science, Osaka University,
Toyonaka 560-0043, Japan

²Department of Physics, Ritsumeikan University, Kusatsu, Shiga 525-8577, Japan
e-mail: asano@vega.ess.sci.osaka-u.ac.jp, fukuyama@se.ritsumei.ac.jp

Received _____; accepted _____

ABSTRACT

Using idealized models of the accretion disk we investigate the relativistic effects on the energy deposition rate via neutrino pair annihilation near the rotation axis of a Kerr black hole. Neutrinos are emitted from the accretion disk. The bending of neutrino trajectories and the redshift due to the disk rotation and gravitation are taken into consideration. The Kerr parameter, a , affects not only neutrinos' behavior but also the inner radius of the accretion disk. When the deposition energy is mainly contributed by the neutrinos coming from the central part, the redshift effect becomes dominant as a becomes large and the energy deposition rate is reduced compared with that neglecting the relativistic effects. On the other hand, for small a the bending effect gets dominant and makes energy increase by factor 2 compared with that neglecting the relativistic effects. For the disk with temperature gradient, the energy deposition rate for a small inner radius of the accretion disk is smaller than that estimated by neglecting the relativistic effects. The relativistic effects, especially for large a , play a negative role in avoiding the baryon contamination problem in gamma-ray bursts.

Subject headings: accretion, accretion disks—black hole physics—gamma rays: bursts

1. INTRODUCTION

The light curves (Fruchter et al. 1999; Harrison et al. 1999; Huang, Dai & Lu 2000) and polarization (Covino et al. 1999; Wijers et al. 1999) of gamma-ray burst afterglows indicate anisotropic central engines. These observations have been considered as evidence that gamma-ray bursts are highly beamed. One of the most probable candidates for the central engine of gamma-ray bursts is the accretion disk around a black hole (Woosley 1993; Popham, Woosley & Fryer 1999; MacFadyen & Woosley 1999; Ruffert & Janka 1999). This system may be formed by the merging of two neutron stars, the merging of a black hole and a neutron star, or the failed supernovae (Woosley 1993).

In these systems the hot accretion disk emits neutrinos and antineutrinos. The neutrino-antineutrino annihilation into electrons and positrons (hereafter neutrino pair annihilation) is a possible energy source of gamma-ray bursts (Paczynski 1990; Mészáros & Rees 1992; Jaroszyński 1996; Janka & Ruffert 1996; Ruffert et al. 1997; Ruffert & Janka 1998, 1999). In order to get the gamma-ray bursts from the relativistic fireball (Shemi & Piran 1990; Rees & Mészáros 1992; Mészáros & Rees 1993; Sari & Piran 1995; Sari, Narayan & Piran 1996), the baryon density in the fireball must be extremely small. Above the accretion disk, the baryon density has the lowest value near the rotation axis. The neutrino pair annihilation in this region has the possibilities of making a clean fireball and of solving the baryon contamination problem. It is important to obtain the energy density profile along the rotation axis in order to discuss the baryon contamination problem and the beaming effect.

The energy deposition rate (hereafter EDR) due to neutrino pair annihilation (Goodman, Dar & Nussinov 1987; Cooperstein, Van Den Horn & Baron 1987; Berezhinsky & Prilutsky 1987) and the gravitational effects on it (Jaroszyński 1993; Ruffert & Janka 1999; Salmonson & Wilson 1999) have been calculated. In the recent study, Asano & Fukuyama (2000; hereafter AF) calculated EDR by neutrinos emitted from the accretion disk. They incorporated the gravitational effects due to the central Schwarzschild black hole and concluded that the gravitational effects do not substantially change EDR. However, AF did not consider the effects of the rotation and temperature gradient of the accretion disk. EDR is proportional to T_{eff}^9 , where T_{eff} is the effective temperature of the accretion disk. Therefore, in order to discuss EDR, we have to include the temperature dependence on radius of the accretion disk.

The relativistic effects consist of three factors: they are the gravitational

redshift, the bending of neutrino trajectories, and the redshift due to the disk rotation. EDR is enhanced by the effect of neutrino bending. However, the redshift due to the disk rotation and gravitation reduces EDR. Thus these effects complicate the estimate of EDR. It is not obvious whether the bending effect becomes dominant and EDR is increased or the redshift effect becomes dominant and EDR is decreased. AF did not consider the Kerr black hole. The rotation of the black hole affects the behavior of neutrinos. In addition the rotation of the black hole makes the inner radius of the accretion disk smaller. These effects may drastically change EDR differently from the case for the Schwarzschild black hole.

In order to discuss quantitatively the central engine of gamma-ray bursts, we need the comprehensive studies of the formation, evolution, temperature dependence, opacity and geometry of the accretion disk and with the mechanism of the energy deposition, its efficiency, the environment around the fireball, the evolution of the fireball and the efficiency of radiation in gamma-ray etc., most of which are affected by the general relativistic effects. We think that these problems will be solved finally by laborious simulations. However, no one reveals the reality of the central engine at the present stage and can say anything affirmative on it. In these situations we discuss qualitatively the problem limiting on the relativistic effects on neutrino pair annihilation which have not been treated seriously in numerical studies. We do not deal with a specific model of gamma-ray bursts, but derive general conclusions on EDR of gamma-ray bursts. Our aim in this work is not to estimate quantitatively the energy of gamma-ray bursts but to study semi-analytically the relativistic effects on EDR above the accretion disk around a rotating black hole. For the above purpose we consider the most idealized situations and compare the results under the same conditions except for the presence and absence of gravitation. We calculate EDR along the rotation axis where the baryon contamination is minimum. The rotation and temperature gradient of the accretion disk are taken into consideration. We assume the most simple models of temperature gradient.

This paper is organized as follows. In section two we formulate the algorithm of EDR calculation along the rotation axis of a Kerr black hole. The results caused by neutrinos emitted from the isothermal disk and from the disk with $T_{\text{eff}} \propto R^{-1}$ and $R^{-0.5}$ are given in section three. The last section is devoted to conclusions.

2. FORMULATION

The formulation developed in this section is obtained by modifying AF's work for the Schwarzschild black hole. In the present case, we consider the following situations. Around a rotating black hole, a hot thin accretion disk is formed and emits a vast number of neutrinos. Because we do not consider a specific model on the central engine of gamma-ray bursts, we consider the most idealized situations irrelevant to detailed models of the disk formation and so on. The inner (outer) edge of this accretion disk is denoted by R_{in} (R_{out}). The geometrically thin disk is assumed to be sufficiently opaque to neutrinos over the whole region. We assume that the self-gravitational effects of the accretion disk is negligible and neutrinos are gravitationally affected only by the central black hole. Of course, these idealizations may be far from the case of the realistic accretion disk. However, we consider that this simple model is sufficient for qualitatively studying the gravitational effects on EDR. The metric around the rotating black hole is given by

$$ds^2 = g_{ij}dx^i dx^j = \left(1 - \frac{r_g r}{\Sigma}\right) c^2 dt^2 - \frac{\Sigma}{\Delta} dr^2 - \left(\frac{r_g r}{\Sigma} a^2 \sin^2 \theta + r^2 + a^2\right) \sin^2 \theta d\varphi^2 - \Sigma d\theta^2 + 2 \frac{r_g r}{\Sigma} a \sin^2 \theta c dt d\varphi. \quad (1)$$

Here $\Sigma \equiv r^2 + a^2 \cos^2 \theta$, $\Delta \equiv r^2 + a^2 - r_g r$ and $r_g = 2GM/c^2$ is the Schwarzschild radius. Also a is the Kerr parameter and takes a value from 0 to $r_g/2$. In this field the eikonal for a massless particle is written as (Landau & Lifshitz 1979)

$$\psi = -\omega_0 t + L\varphi + \psi_r(r) + \psi_\theta(\theta), \quad (2)$$

where ω_0 and L are constants and connected with the energy and the angular momentum around the rotation axis of the black hole, respectively. $\psi_r(r)$ and $\psi_\theta(\theta)$ satisfy the equations

$$\left(\frac{\partial \psi_r(r)}{\partial r}\right)^2 = \frac{1}{\Delta^2} \left[\frac{\omega_0}{c}(r^2 + a^2) - aL\right]^2 - \frac{J^2}{\Delta}, \quad (3)$$

$$\left(\frac{\partial \psi_\theta(\theta)}{\partial \theta}\right)^2 = J^2 - \left[\frac{a\omega_0}{c} \sin \theta - \frac{L}{\sin \theta}\right]^2, \quad (4)$$

where J is a constant (the separation parameter) and connected with the total angular momentum. From equation (2), we obtain the momentum of a neutrino by $p_i = \hbar \partial \psi / \partial x^i$.

As was mentioned in Introduction, we are most interested in EDR via neutrino pair annihilation near the rotation axis, $\theta = 0$. The region of $\theta = 0$ allows analytical treatment because of its symmetric property. In the absence of gravitation, AF indicated that the θ -dependence of EDR is weak for small θ . Although the θ -dependence of the gravitational effects may not necessarily be small, we calculate EDR at $\theta = 0$ and take it as a rough standard in the vicinity of the rotation axis.

It should be remarked that $L = 0$ is required for neutrinos to reach to $\theta = 0$. The eikonal equation only is not sufficient to obtain φ_ν , which is the angle from which meridian the particle is approaching the rotation axis. Here and henceforth the subscript ν and $\bar{\nu}$ denote the quantities related to neutrinos and antineutrinos, respectively. Taking this φ_ν into consideration, the inner product of the momenta, p_ν and $p_{\bar{\nu}}$, at $\theta = 0$ becomes

$$(p_\nu \cdot p_{\bar{\nu}}) = \frac{\varepsilon_\nu \varepsilon_{\bar{\nu}}}{c^2} (1 - \sin \theta_\nu \sin \theta_{\bar{\nu}} \cos(\varphi_\nu - \varphi_{\bar{\nu}}) - \cos \theta_\nu \cos \theta_{\bar{\nu}}). \quad (5)$$

Here the proper energy of neutrinos has been written as

$$\varepsilon_\nu = \hbar \omega_{0\nu} \sqrt{\frac{r^2 + a^2}{\Delta}} = \varepsilon_{0\nu} \sqrt{\frac{r^2 + a^2}{\Delta}}, \quad (6)$$

where $\varepsilon_{0\nu}$ is the energy observed at infinity. The factor $\sqrt{(r^2 + a^2)/\Delta}$ denotes the redshift. θ_ν is defined by

$$\sin \theta_\nu = \frac{\rho_\nu \sqrt{\Delta}}{r^2 + a^2}, \quad (7)$$

with

$$\rho_\nu \equiv \frac{c J_\nu}{\omega_{0\nu}}. \quad (8)$$

A neutrino is emitted from the disk at $(r, \theta) = (R, \pi/2)$, and it arrives at a point $(r, 0)$. Then φ_ν can take values from 0 to 2π at $(r, \theta) = (r, 0)$ from the symmetric property. On the other hand, a neutrino coming from R_{in} forms θ_{m} , the minimum value of θ_ν , and that from R_{out} forms θ_{M} , the maximum value of θ_ν . These values are derived from equation (7) and ρ_ν which is obtained by solving the trajectory equation ($\partial\psi/\partial J^2 = \text{const.}$),

$$\int_0^{\pi/2} \frac{d\theta}{\sqrt{1 - (a/\rho_\nu)^2 \sin^2 \theta}} = \int_C \frac{dr'}{r' \sqrt{(r'/\rho_\nu)^2 (1 + (a/r')^2)^2 - (1 - r_g/r' + (a/r')^2)}}. \quad (9)$$

Here, in the case in which a neutrino passes through the nearest distance, r_0 , before it arrives at $\theta = 0$, the integration for r' is performed from r_0 to R and r . When r' varies monotonically, the integration is performed from the smaller to the larger of r and R . r_0 and ρ_ν are related by

$$\rho_\nu = \frac{r_0(1 + (a/r_0)^2)}{\sqrt{1 - r_g/r_0 + (a/r_0)^2}}. \quad (10)$$

As was derived in AF, EDR via neutrino pair annihilation for a distant observer is expressed as

$$\frac{dE_0(r)}{dt dV} = \frac{21\pi^4}{4} \zeta(5) \frac{K G_F^2}{h^6 c^5} k^9 T_{\text{eff}}^9 (3r_g) F(r), \quad (11)$$

where

$$F(r) = \frac{1}{T_{\text{eff}}^9(3r_g)} \left(\frac{r^2 + a^2}{\Delta} \right)^4 \int_{\theta_m}^{\theta_M} d\theta_\nu \sin \theta_\nu \int_{\theta_m}^{\theta_M} d\theta_{\bar{\nu}} \sin \theta_{\bar{\nu}} \int_0^{2\pi} d\varphi_\nu \int_0^{2\pi} d\varphi_{\bar{\nu}} \\ \times T_0^5(R_\nu) T_0^4(R_{\bar{\nu}}) [1 - \sin \theta_\nu \sin \theta_{\bar{\nu}} \cos(\varphi_\nu - \varphi_{\bar{\nu}}) - \cos \theta_\nu \cos \theta_{\bar{\nu}}]^2. \quad (12)$$

Here the following notations and remarks are in order. By the temperature at $R = 3r_g$, we have represented the effective temperature of the disk T_{eff} which is observed in the comoving frame. T_0 is the temperature of the disk observed at infinity. $dt dV$ denotes $\sqrt{-g} d^4x$. The dimensionless parameter K is 0.12 for ν_e and 0.027 for ν_μ and ν_τ . The Fermi constant G_F^2 is $5.29 \times 10^{-44} \text{cm}^2 \text{MeV}^{-2}$. k and h are the Boltzmann and the Planck constants, respectively. The factor of the redshift in equation (12) and the equation to obtain θ_m and θ_M are different from those for the Schwarzschild metric discussed in AF. Furthermore, although it was assumed in AF that T_0 does not depend on R , we have here assumed that T_0 depends on R . From an angle θ_ν at a point $(r, 0)$, we can trace back a trajectory of a neutrino to the emitted position R_ν on the disk. Thus R_ν and, therefore, $T_0(R_\nu)$ depend on θ_ν , and T_0 is involved in the integrand of θ_ν . This θ_ν -dependence of T_0 is obtained by the numerical calculation of equation (9).

In the locally nonrotating frame (Bardeen, Press, & Teukolsky 1972), a neutrino with $L = 0$ moves normally to the direction of the disk motion. In that case, the temperature suffers the redshift due to the rotation of the disk and gravitation. Namely,

$$T_0(R) = \frac{T_{\text{eff}}(R)}{\gamma} \sqrt{g_{tt} - \frac{g_{t\varphi}^2}{g_{\varphi\varphi}}} = \frac{T_{\text{eff}}(R)}{\gamma} \sqrt{\frac{R^2 + a^2 - r_g R}{R^2 + a^2 + (r_g/R)a^2}}, \quad (13)$$

where $\gamma \equiv 1/\sqrt{1 - v^2/c^2}$ and

$$\frac{v^2}{c^2} = \frac{r_g R (R^2 + a^2 - a\sqrt{2r_g R})^2}{2(R^2 + a\sqrt{r_g R/2})^2 (R^2 + a^2 - r_g R)}, \quad (14)$$

(Chandrasekhar 1983).

Integrating it over φ_ν and $\varphi_{\bar{\nu}}$, $F(r)$ becomes

$$\begin{aligned} F(r) = & \frac{2\pi^2}{T_{\text{eff}}^9(3r_g)} \left(\frac{r^2 + a^2}{\Delta} \right)^4 \left[2 \int_{\theta_m}^{\theta_M} d\theta_\nu T_0^5(\theta_\nu) \sin \theta_\nu \int_{\theta_m}^{\theta_M} d\theta_{\bar{\nu}} T_0^4(\theta_{\bar{\nu}}) \sin \theta_{\bar{\nu}} \right. \\ & + \int_{\theta_m}^{\theta_M} d\theta_\nu T_0^5(\theta_\nu) \sin^3 \theta_\nu \int_{\theta_m}^{\theta_M} d\theta_{\bar{\nu}} T_0^4(\theta_{\bar{\nu}}) \sin^3 \theta_{\bar{\nu}} \\ & + 2 \int_{\theta_m}^{\theta_M} d\theta_\nu T_0^5(\theta_\nu) \cos^2 \theta_\nu \sin \theta_\nu \int_{\theta_m}^{\theta_M} d\theta_{\bar{\nu}} T_0^4(\theta_{\bar{\nu}}) \cos^2 \theta_{\bar{\nu}} \sin \theta_{\bar{\nu}} \\ & \left. - 4 \int_{\theta_m}^{\theta_M} d\theta_\nu T_0^5(\theta_\nu) \cos \theta_\nu \sin \theta_\nu \int_{\theta_m}^{\theta_M} d\theta_{\bar{\nu}} T_0^4(\theta_{\bar{\nu}}) \cos \theta_{\bar{\nu}} \sin \theta_{\bar{\nu}} \right]. \quad (15) \end{aligned}$$

We ignore the rate of deposition energy reabsorbed by the central black hole.

As was estimated in AF, the effect of reabsorption is not so large.

3. RESULTS

3.1. Isothermal Disk

We estimate EDR by neutrinos emitted from the isothermal accretion disk. For this case, neglecting the rotation of the disk and the gravitation of the central black hole, AF calculated EDR in a wide region apart from the rotation axis as well as along it. For the reference, EDR by ν_e over $r = 1.5\text{--}10r_g$ and $\theta \leq \pi/4$ is

$$\frac{dE_0}{dt} = 5.28 \times 10^{52} \left(\frac{kT_{\text{eff}}(3r_g)}{10\text{MeV}} \right)^9 \left(\frac{r_g}{10\text{km}} \right)^3 \text{ ergs s}^{-1}, \quad (16)$$

where $R_{\text{in}} = 3r_g$ and $R_{\text{out}} = 10r_g$. In this case the energy fraction deposited in this region is

$$\frac{\dot{E}_0}{\mathcal{L}} = 2.03 \times 10^{-2} \left(\frac{kT_{\text{eff}}(3r_g)}{10\text{MeV}} \right)^5 \left(\frac{r_g}{10\text{km}} \right), \quad (17)$$

where $\mathcal{L}$ is the neutrino and antineutrino luminosity. If $kT_{\text{eff}}(3r_g)$ is 10 MeV, about 2% of the neutrino energy is deposited in this region. This temperature

is almost the same as the typical temperature in the simulation of neutron star mergers by Ruffert & Janka (1999). We have neglected the decrease of neutrino density due to the annihilation in our formulation. Thus the energy fraction gets larger than one if we adopt a higher temperature than $kT_{\text{eff}}(3r_g) \simeq 20$ MeV in equation (17). This implies that neutrinos become optically thick for the pair annihilation in this case.

In the following, we calculate EDR incorporating the disk rotation and the gravitation of the central black hole. As a matter of convenience, AF assumed that the temperature of the accretion disk is relativistically isothermal, namely, T_0 is constant. In that case, we can estimate equation (11) by calculating the only two neutrino trajectories coming from R_{in} and R_{out} . T_0 goes out of the integral in equation (15). However, it may be more natural to take the effective temperature $T_{\text{eff}}(R)$ as a base since the temperature (and other thermodynamic quantities) is well defined in the comoving coordinates. Thus in this subsection we assume the constant effective temperature, $T_{\text{eff}}(R) = T_{\text{eff}}(3r_g)$. In this case T_0 becomes hotter as R goes to outer region. If we assume $T_{\text{eff}}(3r_g)$ is common, EDR for the constant T_{eff} becomes larger than that for the constant T_0 .

Integration of equation (11) over the volume $dV = (r^2 + a^2 \cos^2 \theta) \sin \theta dr d\theta d\varphi$ gives the energy deposition per unit world time for a distant observer. However, we estimate EDR within the infinitesimal angle $d\theta$ along the rotation axis over $r = 1.5\text{--}10r_g$. We do not consider EDR at $r < 1.5r_g$ since in this region the baryon contamination occurs severely and the reabsorption rate of the deposited energy to the black hole is large. Then EDR is proportional to the dimensionless integral of $G(r) \equiv F(r)(r^2 + a^2)/r_g^2$ over $\hat{r} \equiv r/r_g$;

$$\frac{dE_0}{dt} \simeq 4.41 \times 10^{48} \left(\frac{d\theta}{10^\circ} \right)^2 \left(\frac{kT_{\text{eff}}(3r_g)}{10\text{MeV}} \right)^9 \left(\frac{r_g}{10\text{km}} \right)^3 \int_{1.5}^{10} G(\hat{r}) d\hat{r} \quad \text{ergs s}^{-1}, \quad (18)$$

for ν_e .

We discuss the cases that the dimensionless quantity $a_* \equiv a/0.5r_g$ takes the values 0, 0.9, and 0.99. We adopt the innermost stable orbit as the inner edge of the disk, R_{in} . As a_* takes a larger value, R_{in} becomes smaller. R_{in} are 3, 1.16, and $0.73r_g$ for $a_* = 0, 0.9$, and 0.99 , respectively. For $a_* = 0.99$, R_{in} is inside of the ergosphere. R_{out} is fixed at $10r_g$ in every case.

We plot $G(r)$ in Figure 1 for all these cases. This figure shows that EDR in the case of $a_* = 0$, the Schwarzschild black hole, is enhanced in comparison with the case ignoring the relativistic effects. The disk temperature observed at infinity is higher at the outer region unlike the model of constant T_0 discussed in AF. Correspondingly EDR should be larger than the case of AF. On the other

hand, the Doppler effect due to the rotating disk reduces EDR. Consequently the result is not so different from that of AF.

In Table 1 we list the integral of $G(r)$ over $r = 1.5\text{--}10r_g$. When $R_{\text{in}} = 3r_g$ and the relativistic effects are neglected, electron neutrinos deposit in this region the energy,

$$\frac{dE_0}{dt} \simeq 2.23 \times 10^{51} \left(\frac{d\theta}{10^\circ} \right)^2 \left(\frac{kT_{\text{eff}}(3r_g)}{10\text{MeV}} \right)^9 \left(\frac{r_g}{10\text{km}} \right)^3 \text{ ergs s}^{-1}. \quad (19)$$

The values listed in Table 1 are normalized by this EDR. For $kT_{\text{eff}}(3r_g) = 10$ MeV in this case, the energy deposited in this region is 0.086 % of the neutrino luminosity. As is understood from Table 1 and Figure 1, EDR becomes larger as a_* takes a larger value. The smaller R_{in} causes the larger neutrino luminosity. Although the luminosity observed at infinity for $a_* = 0.9$ is at most 10 % larger than the case for $a_* = 0$, EDR for $a_* = 0.9$ becomes 1.3 times of the latter. For $a_* \geq 0.99$ EDR changes little and practically remains constant. This is because the disk surface does not change very much and the neutrinos emitted from the innermost region suffer the large redshift due to the disk rotation and gravitation for $a_* \geq 0.99$. As is seen from Figure 1, the peak of $G(r)$ approaches to the origin as a_* becomes larger. This is not desirable in order to avoid the baryon contamination problem. However, this negative property is overcome by the increase of the deposited energy.

From the above discussion, we conclude that the relativistic effects make increase EDR along the rotation axis by about factor two, irrespective of a_* . However, the order of EDR is unchanged. Therefore, even if we calculate EDR neglecting the relativistic effects, we can estimate roughly the energy of gamma-ray bursts.

3.2. Disk with Temperature Gradient

EDR via neutrino pair annihilation is strongly dependent on the disk temperature as $dE_0/dt \propto T_{\text{eff}}^9$. Thus the isothermal disk is too crude even for the zeroth-order approximation. In this subsection, we discuss the pair annihilation of neutrinos emitted from the disk with temperature gradient. Here, as before, we do not consider the detailed structure and the formation process of the disk. We have no definite theory on the temperature gradient of the accretion disk as the source of gamma-ray bursts. Here we adopt $T_{\text{eff}}(R) = T_{\text{eff}}(3r_g) \cdot 3r_g/R \propto R^{-1}$ as the simplest and acceptable model (see e.g. Kato, Fukue & Mineshige 1998).

In this model the temperature profiles for different a_* remain common outside of $R = 3r_g$. Actual accretion disks may not be opaque and not hot in the central region. Here, however, we consider the idealized disk model mentioned above. This disk model has the highest temperature at R_{in} where we need the most detailed survey of the gravitational effects. The efficiency of the energy deposition is very high because the innermost temperature is highest.

In this case it is not suitable unlike the isothermal case to discuss based on the disk of $R_{\text{in}} = 3r_g$. As R_{in} gets decreased, the neutrino luminosity becomes very large due to neutrinos coming from the hotter region inside of $R = 3r_g$ and enhances EDR drastically. For instance, in the case ignoring the gravitation and disk rotation, EDR in the region $r = 1.5-10r_g$ and $\theta \leq \pi/4$ is

$$\frac{dE_0}{dt} = 5.76 \times 10^{50} \left(\frac{kT_{\text{eff}}(3r_g)}{10\text{MeV}} \right)^9 \left(\frac{r_g}{10\text{km}} \right)^3 \text{ ergs s}^{-1} \text{ for } R_{\text{in}} = 3r_g, \quad (20)$$

$$= 1.05 \times 10^{53} \left(\frac{kT_{\text{eff}}(3r_g)}{10\text{MeV}} \right)^9 \left(\frac{r_g}{10\text{km}} \right)^3 \text{ ergs s}^{-1} \text{ for } R_{\text{in}} = 1.16r_g \quad (21)$$

where we have adopted $R_{\text{out}} = 10r_g$ as before. These values have been calculated by use of the formulation in AF, taking the temperature gradient into consideration. For $R_{\text{in}} = 3r_g$, in comparison with the isothermal case (see equation (16)), the luminosity and EDR are reduced roughly by factor 10 and 100, respectively for the same $T_{\text{eff}}(3r_g)$. Therefore, EDR gets less efficient. For $R_{\text{in}} = 1.16r_g$, on the other hand, the luminosity is about 0.7 times that of the isothermal disk with $R_{\text{in}} = 3r_g$. However, EDR becomes about two times larger and the efficiency is improved compared with the isothermal disk for the same $T_{\text{eff}}(3r_g)$. We have listed in Table 2 EDR along the rotation axis over $r = 1.5-10r_g$ for various R_{in} in the case ignoring the relativistic effects. $R_{\text{in}} = 1.45$ and 0.59 correspond to the innermost stable orbit for $a_* = 0.8$ and 0.999 , respectively. Table 2 shows that EDR increases rapidly due to the decrease of R_{in} . The neutrino luminosity emitted from the disk with $R_{\text{in}} = 0.73r_g$ becomes only 19 times that of the case for $R_{\text{in}} = 3r_g$, whereas EDR of the former case increases by factor 1400 compared with the latter one.

In the following we estimate EDR taking into consideration the gravitation of the central black hole and the disk rotation. The relativistic effects are estimated comparing with EDR for the same R_{in} without the relativistic effects. We adopt $R_{\text{out}} = 10r_g$ as before. We plot $G(r)$ in Figure 2 for the Schwarzschild black hole. In the case of $T_{\text{eff}} \propto R^{-1}$, EDR decreases rapidly with r unlike the isothermal case. EDR increases at small r due to the gravitational lensing. Since the neutrino energy decreases due to the gravitational redshift, EDR is

less efficient at r larger than $r = 6r_g$ than that without the relativistic effects. As in the isothermal case, the relativistic effects make the point of the maximum rate approach to the origin. As is shown in Table 3, the relativistic effects for $a_* = 0$ enhance EDR by factor 1.7. This is the same tendency as in the case of the isothermal disk.

Whereas, for $a_* = 0.8$ it goes from Figure 3 that the relativistic effects enhance EDR only in the very neighborhood of the black hole and reduce it at distant places due to the redshift. These relativistic effects are disadvantageous to avoid the baryon contamination by the energy deposition at distant places from the central object. The peak of EDR lies in $r < 1.5r_g$. Table 3 indicates that EDR for $a_* = 0.8$ over $r = 1.5-10r_g$ is not changed very much compared with the case ignoring the relativistic effects.

As a_* takes a larger value, EDR itself increases. For $a_* \geq 0.9$, however, the energy reduction by the relativistic effects becomes larger as a_* increases (see Table 3). For $a_* = 0.9$, as is shown in Figure 4, $G(r)$ takes lower values than those in the case ignoring the relativistic effects in almost all regions and we cannot expect a large deposition at distant places from the center. In Figure 5, $G(r)$ for $a_* = 0.99$ is much smaller than that ignoring the relativistic effects in all region, especially at distant places. These results are disadvantageous to avoid the baryon contamination problem in gamma-ray bursts. In these cases, EDR is largely owing to the neutrinos emitted from the innermost hot region of the disk. These neutrinos suffer the large redshift due to the rapid disk rotation and gravitation and cannot effectively deposit energy at $r > R$. Figure 5 clearly shows that EDR substantially increases when R_{in} decreases in the case neglecting the relativistic effects. On the other hand, if we plot $G(r)$ for $a_* = 0.999$ in Figure 5 taking into consideration the relativistic effects, $G(r)$ overlaps almost completely with those for $a_* = 0.99$ with the relativistic effects (solid line).

The neutrino luminosity observed at infinity for, as an example, $a_* = 0.99$ is reduced to 0.24 times that ignoring the relativistic effects. Thus the redshift seriously affects the luminosity, which results in Table 3. However, the results of Table 3 cannot be explained only by the change of the luminosity. For when a_* goes from 0.99 to 0.999, the luminosity increases by factor 1.2 but EDR remains almost constant. Whereas if we neglect the relativistic effects, the luminosity and EDR increase, respectively, by factor 1.5 and 2 for the same change of R_{in} . The different pattern of the EDR growth between these two cases indicates that the relativistic effects on the reaction rate of neutrino pair annihilation are also important. Even if a_* changes, EDR at distant places is practically unchanged

for $a_* > 0.8$. Most of the energy growth due to the increase of a_* occurs near the center.

As mentioned above, when a_* goes beyond 0.99, EDR itself remains almost constant. For $T_{\text{eff}} \propto R^{-1}$, even if a_* approaches to 1 beyond 0.99, the deposited energy does not increase because of the redshift due to the disk rotation and gravitation unlike the case neglecting the gravitational effects. The obtained EDR is fairly small compared with that in the case where R_{in} approaches to $0.5r_g$ and the relativistic effects are neglected. Namely, as a_* approaches to one, the value of about 2.7 times equation (19) is the rough maximum value of EDR in this model. The fraction of EDR against the neutrino luminosity increases as a_* becomes large. For $a_* = 0.99$, if we take account of the relativistic effects, the neutrino luminosity observed at infinity reduces by factor 0.4 compared with the case of equation (19). Therefore, the efficiency of EDR becomes 6.5 times that for the case of equation (19). For the same $T_{\text{eff}}(3r_g)$, Figure 1 and 5 indicate that EDR at distant places above the disk with $T_{\text{eff}} \propto R^{-1}$ is much smaller than those above the isothermal disk, although the integrated deposition rates are not so different. Thus the isothermal disk is more advantageous than the disk with $T_{\text{eff}} \propto R^{-1}$ to avoid the baryon contamination problem.

We have calculated for the two extreme disk models, $T_{\text{eff}} \propto R^0$ and R^{-1} , on behalf of various disk models. The calculations for $T_{\text{eff}} \propto R^{-1}$ show the following things: If the deposited energy is mainly contributed by the neutrinos coming from the central region, the bending effect which increases EDR is dominant for large R_{in} , and the redshift effect which decrease EDR is dominant for small R_{in} . Namely, as R_{in} approaches the horizon, the redshift effect overwhelms the bending effect. If we assume a model with the more flat temperature gradient than $T_{\text{eff}} \propto R^{-1}$ (the numerical work of Popham, Woosley & Fryer (1999) predicts such disks), the contribution of the neutrinos coming from central region is relatively diminished compared with the case of $T_{\text{eff}} \propto R^{-1}$. Therefore the increase of EDR due to the decrease of R_{in} is not so prominent as in the cases of Table 2 and 3, and correspondingly the relativistic effects are not so manifest as shown in Table 3. However, the qualitative result that the redshift effect becomes dominant according to the decrease of R_{in} (or according to the growth of a_*) is unchanged. We have listed the relativistic effects for the disk with $T_{\text{eff}}(R) = T_{\text{eff}}(3r_g) \cdot \sqrt{3r_g/R} \propto R^{-0.5}$ in Table 4. This table strongly supports our consideration mentioned above. Since the redshift effect becomes dominant as a_* increases, the enhancement factor of EDR due to the relativistic effects has the maximum value in the Schwarzschild case. The enhancement factor in Table 4 is 1.7, which is the same as those in the former two disk

models. From these arguments it is conjectured that the enhancement factor due to the relativistic effects probably does not exceed 2 in the disk models with simple temperature dependences. Also in Table 4, for $a_* > 0.99$, EDR itself remains almost constant as in the other two cases.

We are not able to investigate all possible models with different temperature distributions. In order to study the qualitative properties of relativistic effects we have calculated only the simple models mentioned above. However, if the opacity and the density in the central region are low, EDR is not dominated by the neutrinos coming from that region. The redshift effect is not so prominent as that of our cases. In that case the role of R_{in} is replaced by the radius at which T_{eff} has the highest value. As this radius moves outward, the bending effect gradually becomes dominant and the enhancement factor due to the relativistic effects may approach 1.7 of $a_* = 0$ case.

4. CONCLUSIONS

In this article we have investigated the relativistic effects on EDR via neutrino pair annihilation near the rotation axis of a Kerr black hole with the thin accretion disk. The energy deposited over $r = 1.5\text{--}10r_g$ is our concern in this paper. The bending of neutrino trajectories and the redshift due to the disk rotation and gravitation have been taken into consideration. The Kerr parameter, a , affects not only neutrinos' behavior but also the inner radius of the accretion disk. If the accretion disk is isothermal, the relativistic effects make increase EDR by about factor two, irrespective of a .

On the other hand, if the temperature of the accretion disk behaves as $T_{\text{eff}} \propto R^{-1}$, the increase of a enhances EDR drastically. However, the most of the pair annihilation occur near the central black hole. This is disadvantageous to avoid the baryon contamination problem in gamma-ray bursts. When a_* goes beyond 0.99, EDR remains almost constant unlike the case without the relativistic effects and is much smaller than EDR in the case where R_{in} approaches to $0.5r_g$ and the relativistic effects are neglected. This is because the neutrinos, emitted from the innermost hot spot of the disk, suffer the large redshift. This is the case also for the disk with the more flat temperature gradient and for the isothermal disk. Therefore, these results indicate that EDR hardly increases, irrelevant to models of the disk temperature, even if we consider the disk with small R_{in} .

From the above arguments the qualitative properties of the relativistic effects on neutrino pair annihilation are summarized as follows. When the deposition energy is mainly contributed by the neutrinos coming from the central part, the redshift effect becomes dominant as this active part approaches to the horizon and EDR is reduced compared with that neglecting the relativistic effects. On the other hand, as the part emitting the dominant neutrinos goes away from the horizon, the bending effect gets dominant to make EDR increase by factor 2 compared with that neglecting the relativistic effects.

These qualitative conclusions may be helpful for numerical simulations of gamma-ray burst sources which neglect the relativistic effects in propagation of neutrinos. In order to discuss quantitatively the central engine of gamma-ray bursts, one needs numerical simulations and several groups are acting in this field (e.g. Popham, Woosley & Fryer 1999; MacFadyen & Woosley 1999; Ruffert & Janka 1999). The gravitational effects on neutrino pair annihilation may quantitatively change the results of such simulations. As it comes from our calculations, the estimated power of a gamma ray burst may be changed by a factor less than two if the temperature gradient is relatively small, and the neutrinos emitted close to the inner adge of the disk do not dominate. However, if a_* is close to one and EDR is strongly dominated by the neutrinos coming from the central region, one can not give a reliable result without taking the redshift effect into consideration.

We are greatly indebted to F. Takahara for useful discussion. We are also grateful to S. Kobayashi and for helpful advice and comments. This work was partly supported by a Research Fellowship of the Japan Society for the Promotion of Science.

References

- Asano, K., & Fukuyama, T. 2000, *ApJ*, 531, 949 (AF)
- Bardeen, J. M., Press, W. H., & Teukolsky, S. A. 1972, *ApJ*, 178, 347
- Berezinsky, V. S., & Prilutsky, O. F. 1987, *A&A*, 175, 309
- Chandrasekhar, S. 1983, *The Mathematical Theory of Black Holes* (New York: Oxford)
- Cooperstein, J., Van Den Horn, L. J., & Baron, E. 1987, *ApJ*, 321, L129
- Covino, S., et al. 1999, *A&A*, 348, L1
- Fruchter, A. S., et al. 1999, *ApJ*, 519, L13
- Goodman, J., Dar, A., & Nussinov, S. 1987, *ApJ*, 314, L7
- Harrison, F. A., et al. 1999, *ApJ*, 523, L121
- Huang, Y. F., Dai, Z. G., & Lu, T. 2000, *A&A*, 355, L43
- Janka, H.-T., & Ruffert, M. 1996, *A&A*, 307, L33
- Jaroszyński, M. 1993, *ActaAstron*, 43, 183
- Jaroszyński, M. 1996, *A&A*, 305, 839
- Kato, S., Fukue, J., & Mineshige, S. 1998, *Black-hole Accretion Disks* (Kyoto: Kyoto University)
- Landau, L. D., & Lifshitz, E. M. 1979, *Classical Theory of Fields* (London: Pergamon)
- MacFadyen, A., & Woosley, S. E. 1999, *ApJ*, 524, 262
- Mészáros, P., & Rees, M. J. 1992, *MNRAS*, 257, 29p
- Mészáros, P., & Rees, M. J. 1993, *ApJ*, 405, 278]
- Paczynski, B. 1990, *ApJ*, 363, 218
- Popham, R., Woosley, S. E., & Fryer, C. 1999, *ApJ*, 518, 356
- Rees, M. J., & Mészáros, P. 1992, *MNRAS* 258, 41p
- Ruffert, M., & Janka, H.-T. 1998, *A&A*, 338, 535
- Ruffert, M., & Janka, H.-T. 1999, *A&A*, 344, 573
- Ruffert, M., Janka, H.-T., Takahashi, K., & Schäfer, G. 1997, *A&A*, 319, 122
- Salmonson, J. D., & Wilson, J. R. 1999, *ApJ* 517, 859

Sari, R., Narayan, R., & Piran, T. 1996 ApJ, 473, 204

Sari, R., & Piran, T. 1995, ApJ, 455, L143

Shemi, A., & Piran, T. 1990, ApJ, 365, L55

Wijers, R. A. M. J., et al. 1999, ApJ, 523, L33

Woosley, S. E. 1993, ApJ, 405, 273

Figure captions

Fig. 1.— The plot of $G(r)$ (solid line) against r/r_g for the isothermal disk. The dashed line corresponds to the case for $R_{\text{in}} = 3r_g$ ignoring the relativistic effects.

Fig. 2.— The plot of $G(r)$ (solid line) against r/r_g for the case of $a_* = 0$ and $T_{\text{eff}} \propto R^{-1}$. The dashed line corresponds to the case for the same R_{in} ignoring the relativistic effects. Contrary to Figure 1, the vertical axis is measured by the logarithmic scale.

Fig. 3.— Same as Figure 2 but for $a_* = 0.8$.

Fig. 4.— Same as Figure 2 but for $a_* = 0.9$.

Fig. 5.— Same as Figure 2 but for $a_* = 0.99$. The dashed line and dotted line correspond to the cases for $R_{\text{in}} = 0.73$ ($a_* = 0.99$) and $0.59r_g$ ($a_* = 0.999$) ignoring the relativistic effects, respectively.

a_*	0	0.9	0.99
EDR	1.7	2.2	2.3

Table 1. EDR within the infinitesimal angle $d\theta$ along the rotation axis over $r = 1.5\text{--}10r_g$ for the isothermal disk. The values are normalized to unity when $R_{\text{in}} = 3r_g$ and the relativistic effects are neglected.

R_{in}/r_g	3	1.45	1.16	0.73	0.59
EDR	0.011	0.70	2.1	15	31
$\mathcal{L}$	0.09	0.41	0.65	1.7	2.5

Table 2. EDR along the rotation axis over $r = 1.5\text{--}10r_g$ for the disk with $T_{\text{eff}} \propto R^{-1}$ ignoring the relativistic effects. The given values are normalized by the value of equation (19). The neutrino luminosity $\mathcal{L}$ are also listed. Each $\mathcal{L}$ is normalized by the luminosity in the case of equation (19).

a_*	0	0.8	0.9	0.99	0.999
R_{in}/r_g	3	1.45	1.16	0.73	0.59
EDR	1.7 (0.019)	1.1 (0.79)	0.73 (1.5)	0.18 (2.6)	0.087 (2.7)
$\mathcal{L}$	0.62 (0.056)	0.43 (0.18)	0.36 (0.24)	0.24 (0.40)	0.19 (0.47)

Table 3. EDR along the rotation axis over $r = 1.5\text{--}10r_g$ for the disk with $T_{\text{eff}} \propto R^{-1}$ including the relativistic effects. The neutrino luminosities $\mathcal{L}$ observed at infinity are also listed. Each value is divided by that with the same R_{in} ignoring the relativistic effects. The values normalized by the value in the case of equation (19) are also listed as parenthetic values.

a_*	0	0.8	0.9	0.99	0.999
R_{in}/r_g	3	1.45	1.16	0.73	0.59
EDR	1.7 (0.10)	1.4 (0.40)	1.2 (0.47)	0.81 (0.54)	0.68 (0.55)
$\mathcal{L}$	0.69 (0.17)	0.67 (0.23)	0.66 (0.24)	0.64 (0.33)	0.63 (0.35)

Table 4. Same as Table 3 but for $T_{\text{eff}} \propto R^{-0.5}$.









